arXiv:1208.5594v1 [math.CO] 28 Aug 2012

,

LASSOING AND CORALLING ROOTED PHYLOGENETIC TREES

KATHARINA T. HUBER AND ANDREI-ALIN POPESCU

ABSTRACT. The construction of a dendogram on a set of individuals is a key component of a genomewide association study. However even with modern sequencing technologies the distances on the individuals required for the construction of such a structure may not always be reliable making it tempting to exclude them from an analysis. This, in turn, results in an input set for dendogram construction that consists of only partial distance information. By formalizing a dendogram in terms of an edge-weighted, rooted phylogenetic tree on a pre-given finite set X with $|X| \geq 3$ whose edge-weighting is equidistant and a set of partial distances on X in terms of a set $\mathscr{L}$ of 2-subsets of X, we investigate the problem of when such a tree is lassoed, that is, uniquely determined by the elements in $\mathscr{L}$. For this we consider four different formalizations of the idea of "uniquely determining" giving rise to four distinct types of lassos. We present characterizations for all of them in terms of the child-edge graphs of the interior vertices of such a tree. Our characterizations imply in particular that in case the tree in question is binary then all four types of lasso must coincide.

Keywords rooted phylogenetic tree, lasso, corall, partial distance

1. INTRODUCTION

Years of selective breeding have resulted in large numbers of different varieties for, for example, oilseed rape and rice and also numerous animal breeds including dogs and chicken. Genomewide association studies constitute a powerful tool to try and link the observed phenotypic variability between the varieties (we will collectively refer to a variety and a breed as a variety) with variations in the genomes of the variety. A key component of such a study is phenetic clustering whereby one aims to construct a dendogram for a set of individuals from within a variety of interest indicating levels of similarity between them. Using some sort of distance measure this similarity can be based on e. g. morphological traits such as grain type or Single Nucleotide Polymorphism (SNP) markers obtained through next generation sequencing technology (see e. g. Harper et al (in press); Muir et al (2008); Zhang et al (2010) for examples of such studies).

From a formal point of view, a dendogram can be thought of as a pair (T,ω) consisting of (i) a rooted tree T with leaf set a given non-empty finite set X (e. g. individuals), no degree two vertices but possibly a distinguished vertex ρ_T of T called the root of T, and all other non-leaf vertices of T of degree at least three (we will refer to such a tree simply as an *X-tree*), and (ii) an edge-weighting $\omega : E(T) \to \mathbb{R}_{\geq 0}$ for T that is equidistant which means that the induced distance $D_{(T,\omega)}(\rho_T, x)$ from ρ_T to every leaf $x \in X$ of T is the same (as with all relevant concepts, we refer the reader to the next section for a precise definition). With and without the equidistance requirement such pairs (T,ω) have generated a lot of interest in the literature and so it is not surprising that numerous deep results for them are known provided the distance information from which to construct T and ω is complete in the sense that for all elements x and y in X the distance between x and y is given (see e. g. Dress et al (2012a); Semple and Steel (2003)).

Date: August 29, 2012

However even for data generated with modern sequencing technologies the required distance measures need not always be reliable (or may simply be missing) resulting in only partial distance information for dendogram reconstruction. From the perspective of the aforementioned formalization of such a structure, the problem thus becomes (a) how to construct an *equidistant X-tree* (i. e. an X-tree with equidistant edge weighting) from partial distance information and, (b) if such a tree exists, when is it lassoed by that information, that is, unique in the sense that any two equidistant X-trees for which, with regards to the known distances on X, the induced distances on X coincide must be equivalent?

Although approaches for tackling the first problem exist in the form of, for example, an approach introduced in De Soete (1984) not much is known about the second. A notable exception is a study in Dress et al (2012b) carried out for the unrooted analogon of an equidistant X-tree. Viewing partial distance information on a set X as a set of *cords*, that is, subsets of X of size two, the authors considered the following four natural interpretations of the above uniqueness problem. Namely, given a set $\mathscr{L} \subseteq \binom{X}{2}$ and an edge-weighted unrooted phylogenetic tree T on X when

(i) is the edge-weighting of T uniquely determined by $\mathscr{L}$?,
(ii) is the shape i. e. the topology of T uniquely determined by $\mathscr{L}$?,
(iii) are both the edge-weighting and the topology of T uniquely determined by $\mathscr{L}$?, and
(iv) when is the topology of T uniquely determined by $\mathscr{L}$ up to T being obtained from another X-tree by collapsing edges?.

Formalized as $\mathscr{L}$ being an edge-weight/topological/strong/weak lasso for an unrooted phylogenetic tree with leaf set X, the authors of Dress et al (2012b) showed that all four concepts are distinct. Also, they presented results that allowed them to not only investigate the above types of lasso from a recursive point of view but also characterize under what circumstances a specifically constructed set of cords is a topological lasso (see Section 8 for more on this). However a characterization for the general case eluded them.

Replacing the concept of an edge-weight lasso by that of an equidistant lasso to reflect the fact that for the edge-weighted X-trees of interest here the induced distance from the root to any leaf of such a tree is the same, we show that for X-trees the situation changes. More precisely, we present for an X-tree T characterizations for when a set $\mathscr{L} \subseteq \binom{X}{2}$ is weak lasso for T (Theorem 5.1), an equidistant lasso for T (Theorem 6.1), and for when it is a topological lasso for T (Theorem 7.1) in terms of the child edge graph $G_T(\mathscr{L}, v)$ that can be canonically associated to every non-leaf vertex v of T via its child edges. Our characterizations can be thought of as a spectrum on the connectedness of that graph with the extreme situations being an equidistant lasso and a topological lasso. They imply that every topological lasso and every non-empty weak lasso must be an edge-weight lasso ((Corollaries 7.2 and 6.2) and that in case T is binary the notions of an equidistant lasso and a topological lasso (and thus a weak lasso) coincide. Consequently, every edge-weight/topological lasso is also a strong lasso in that case (Corollary 7.2). We also investigate two special types of sets of cords originally introduced in Dress et al (2012b) in the light of our findings above. This investigation shows in particular that it is possible for the concepts of an equidistant and a topological lasso to coincide without the X-tree they are referring to being binary.

The outline of the paper is as follows. In the next section, we introduce relevant terminology. In Section 3, we present first characterizations for when a set of cords is a topological/equidistant/weak lasso for an X-tree (Theorem 3.2). In Section 4, we introduce the child-edge graph associated to a non-leaf vertex of an X-tree and present first

properties of it concerning coralling sets of cords. In Section 5, we establish Theorem 5.1. In Section 6, we show Theorem 6.1 and in Section 7, we prove Theorem 7.1. In Section 8, we present two general ways for constructing for an X-tree T two different sets of cords of $\binom{X}{2}$ and discuss their properties in in the context of lassoing and coralling T. We conclude with Section 9 where we also present some open problems.

2. PRELIMINARIES

Assume from now on that X is always a finite set with at least 3 elements. For a cord of X with elements a and b we write ab rather than $\{a,b\}$.

Suppose for the following that T is an X-tree. Then we call a vertex of T that is not a leaf of T an *interior vertex* of T, an edge e that is incident with a leaf of T a *pendant edge* of T, and an edge of T that is incident with two interior vertices of T an *interior edge* of T. We denote the set of all interior vertices of T by $V^o(T)$ and the root of T by ρ_T. Let T' be a further X-tree. Then we say that T and T' are *equivalent* if there exists a bijection $\phi : V(T) \to V(T')$ that extends to a graph isomorphism between T and T' that is the identity on X and maps the root ρ_T of T to the root $\rho_{T'}$ of T'. Also, we say that T' *refines* T if, up to equivalence, T can be obtained from T' by collapsing edges of T' (see e. g. Semple and Steel (2003)). In that case, we will also call T' a *refinement* of T. Note that T is its own refinement if and only if T is *binary*, that is, every interior vertex of T has degree three but the root which has degree two.

Let ω denote an *edge-weighting for* T, that is, a map $\omega : E(T) \to \mathbb{R}_{\geq 0}$. Then we denote by (T,ω) the edge-weighted X-tree whose underlying X-tree and edge-weighting are T and ω, respectively, and by $D_{(T,\omega)}$ the distance induced by ω on $V(T)$. We call ω *proper* if $\omega(e) > 0$ holds for all interior edges e of T and we call it *equidistant* if

(E1) $D_{(T,\omega)}(x,\rho_T) = D_{(T,\omega)}(y,\rho_T)$, for all $x,y \in X$, and
(E2) $D_{(T,\omega)}(x,u) \geq D_{(T,\omega)}(x,v)$, for all $x \in X$ and any $u,v \in V(T)$ such that u is encountered before v on the path from ρ_T to x.

Note that Property (E2) implies that, for all interior vertices v of T and all leaves $x,y \in X$ of T for which v lies on the path from x to y, we have $D_{(T,\omega)}(x,v) = D_{(T,\omega)}(y,v)$. Also note that our definition of an equidistant edge-weighting is slightly different from the one given in Semple and Steel (2003) in so far that ω is a map into $\mathbb{R}_{\geq 0}$ and not into $\mathbb{R}$, as in Semple and Steel (2003).

Suppose $\mathscr{L} \subseteq \binom{X}{2}$ is a set of cords. If T' is a further X-tree and ω and ω' are proper edge-weightings for T and T', respectively, we say that (T,ω) and (T',ω') are $\mathscr{L}$*-isometric* if $D_{(T,\omega)}(x,y) = D_{(T',\omega')}(x,y)$ holds for all cords $xy \in \mathscr{L}$. Canonically extending the corresponding concepts introduced in Dress et al (2012b) for unrooted phylogenetic trees on X to X-trees, we say that $\mathscr{L}$ is

(i) an *equidistant lasso for* T if, for all equidistant, proper edge-weightings ω and ω' of T, we have that $\omega = \omega'$ holds whenever (T,ω) and (T,ω') are $\mathscr{L}$-isometric
(ii) a *topological lasso for* T if, for every X-tree T' and any equidistant, proper edge-weightings ω of T and ω' of T', respectively, we have that T and T' are equivalent whenever (T,ω) and (T',ω') are $\mathscr{L}$-isometric.
(iii) is a *strong lasso for* T if $\mathscr{L}$ is simultaneously an equidistant and a topological lasso for T.
(iv) a *weak lasso for* T if, for every X-tree T' and any equidistant, proper edge-weightings ω of T and ω' of T', respectively, we have that T is refined by T' whenever (T,ω) and (T',ω') are $\mathscr{L}$-isometric.

Also, we say that a set $\mathscr{L} \subseteq \binom{X}{2}$ of cords is an *equidistant/topological/weak/strong lasso on X* if there exists an X-tree T such that $\mathscr{L}$ is an equidistant/topological/weak/strong lasso for T. For the convenience of the reader, we illustrate the above types of lassos in Fig. 1 for $X = \{a,b,c,d,e\}$.




FIGURE 1. For $X = \{a,b,c,d,e\}$, the set $\mathscr{L} = \{ab,cd,de\}$ is an equidistant lasso for T, the set $\mathscr{L} = \{ab,ac,bc,bd,de\}$ is a topological lasso for T and also a strong lasso for T, and the set $\mathscr{L} = \{ab,bc,cd,de\}$ is a weak lasso for T.

Note that we will also say that a set $\mathscr{L}$ of cords of X *coralls* an X-tree T if $\mathscr{L}$ is a weak lasso for T, that a topological lasso for T is in particular a weak lasso for T, and that the notions of a topological lasso for T and a weak lasso for T coincide if T is binary. Also note that for $\mathscr{L}$ to be a topological/equidistant lasso we must have that $\mathscr{L} \neq \emptyset$. However $\mathscr{L} \neq \emptyset$ need not hold for $\mathscr{L}$ to corall an X-tree as every subset of $\binom{X}{2}$ including the empty-set coralls the *star-tree on X*, that is the tree with a unique interior vertex w and leaf set X.

3. A FIRST CHARACTERIZATION OF A TOPOLOGICAL/WEAK/EQUIDISTANT LASSO

In this section we present a first characterization for when a set of cords of X is a topological/weak/equidistant lasso for an X-tree. To establish this characterization, we require further definitions and notations.

Suppose for the following that T is an X-tree and that $v \in V^o(T)$. Then we call an edge e of T incident with v that is not crossed by the path from the root ρ_T of T to v a *child edge* of v. If e is incident with v but lies on the path from ρ_T to v then we call it a *parent edge* of v. In the former case, we call the vertex incident with that edge but distinct from v a *child* of v and in the latter a *parent* of v. We call a vertex w of T distinct from v a *descendant* of v if there exists a path from v to w (possibly of length one) that crosses a child of v and denote the set of leaves of T that are also descendants of v by $L_T(v)$. If v is a leaf of T, then we put $L_T(v) := \{v\}$. Also if there is no ambiguity as to which X-tree T we are referring then we will write $L(v)$ rather than $L_T(v)$.

Suppose T is an X-tree. Then, for all $x \in X$, we denote the edge of T incident with x by e_x and the parent of x by v_x. Moreover, we call a non-empty subset $L \subsetneq X$ of leaves of T that all have the same parent $v \in V^o(T)$ a *pseudo-cherry* of T if $L = L(v)$. In that case, we also call v the *parent* of that pseudo-cherry. If $\{x_1,\ldots,x_k\}$, $k \geq 1$, is a pseudo-cherry of some X-tree T then we will sometimes write $x_1,\ldots,x_k$ rather than $\{x_1,\ldots,x_k\}$. Note that every X-tree on three or more leaves that is not the star-tree on X must contain at least one pseudo-cherry. Also note that in case $|L| = 2$ the definition of an pseudo-cherry reduces to that of a cherry in the usual sense (see e.g. Semple and Steel (2003)). In the special case that $|X| = 3$, say $X = \{a,b,c\}$, and that T has a cherry, a,b say, we call T a *triplet* and denote T by $ab|c$ (or, equivalently, $c|ab$).

Now assume again that T is an arbitrary X-tree. For any non-empty subset $Y \subseteq X$, we denote by $T|_Y$ the minimum spanning tree in T that connects the leaves in Y (where we *suppress* resulting interior vertices v of T_Y with a single child, that is, if the parent of v is w and u is the unique child of v then we delete v plus its incident edges from $T|_Y$ and add the edge $\{w,u\}$ to $E(T|_Y)$). Note that T_Y is clearly a Y-tree. We say for three pairwise distinct elements $a,b,c \in X$ that T *displays* the triplet $ab|c$ if $T|_{\{a,b,c\}}$ is equivalent with $ab|c$. We denote by $\mathscr{R}(T)$ the set of triplets displayed by T. As is well-known, any X-tree T can display at most $\binom{|X|}{3}$ triplets with equality holding if and only if T is binary. Furthermore, any X-tree T is uniquely determined by the set $\mathscr{R}(T)$ in the sense that if T' is a further X-tree and $\mathscr{R}(T) = \mathscr{R}(T')$ holds then T and T' must be equivalent (see e. g. (Dress et al, 2012a, Chapter 9) and (Semple and Steel, 2003, Section 6.4)).

Observe that if T is an X-tree, ω is an equidistant, proper edge-weighting for T, and $a,a',b \in X$ are pairwise distinct elements then either all three pairwise distances induced on $Z := \{a,a',b\}$ must coincide or two of the distances induced by it must coincide and the third one must be strictly less than that distance. Moreover, $D_{(T,\omega)}(a,a') < D_{(T,\omega)}(a,b) = D_{(T,\omega)}(a',b)$ holds if and only if $aa'|b \in \mathscr{R}(T)$ and we have $D_{(T,\omega)}(a,a') = D_{(T,\omega)}(a,b) = D_{(T,\omega)}(a',b)$ if and only if the Z-tree $T|_Z$ is the star-tree on Z.

The next result is fundamental for the paper.

Lemma 3.1. *Suppose that T and T' are two X-trees with equidistant, proper edge-weightings ω and ω', respectively, and let $a,a',b \in X$ denote three pairwise distinct elements such that $D_{(T,\omega)}(a,a') = D_{(T',\omega')}(a,a')$ and $D_{(T,\omega)}(a,b) = D_{(T',\omega')}(a,b)$. Then the following hold:*

(i) *If $D_{(T,\omega)}(a,a') < D_{(T,\omega)}(a,b) = D_{(T,\omega)}(a',b)$ then*

$$D_{(T',\omega')}(a,a') < D_{(T',\omega')}(a,b) = D_{(T',\omega')}(a',b). \tag{1}$$

In particular, $D_{(T',\omega')}(a',b) = D_{(T,\omega)}(a',b)$.

(ii) *$aa'|b \in \mathscr{R}(T)$ if and only if $aa'|b \in \mathscr{R}(T')$.*

(iii) *If $D_{(T,\omega)}(a',b) = D_{(T',\omega')}(a',b)$ then $T|_Z$ is the star-tree on $Z := \{a,a',b\}$ if and only if $T'|_Z$ is the star-tree on Z.*

Proof. (i): Note that $D_{(T,\omega)}(a,a') = D_{(T',\omega')}(a,a')$ combined with $D_{(T,\omega)}(a,b) = D_{(T',\omega')}(a,b)$ implies that we cannot have $D_{(T',\omega')}(a,a') = D_{(T',\omega')}(a,b)$ as this would imply that $D_{(T,\omega)}(a,a') = D_{(T,\omega)}(a,b)$ which is impossible. But then our assumptions imply that $D_{(T',\omega')}(a,a') = D_{(T',\omega')}(a',b)$ cannot hold either. Inequality (1) now follows from the observation preceding the statement of the lemma. In particular, this implies $D_{(T',\omega')}(a',b) = D_{(T',\omega')}(a,b) = D_{(T,\omega)}(a,b) = D_{(T,\omega)}(a',b)$.

(ii) & (iii): This is an immediate consequence of (i) and the observation preceding the statement of the lemma. □

To be able to state the next result we require a further definition. Suppose $\mathscr{L} \subseteq \binom{X}{2}$ and T is an X-tree. Let $x,y \in X$ be two distinct leaves of T that are contained in the same a pseudo-cherry of T. Then we put

$$\mathscr{L}_1(x,y) := \{ab \in \mathscr{L} : x \notin \{a,b\}\} \cup \{ay : ax \in \mathscr{L}\}.$$

Note that $\mathscr{L} = \emptyset$ if and only if $\mathscr{L}_1(x,y) = \emptyset$.

Theorem 3.2. *Suppose that $\mathscr{L} \subseteq \binom{X}{2}$ is a set of cords and that T is an X-tree. Let $x,y \in X$ denote two distinct leaves of T that are contained in the same pseudo-cherry of T. Then the following hold.*

(i) *$\mathscr{L}$ is a topological lasso for T if and only if $\mathscr{L}_1(x,y)\cup\{xy\}$ is a topological lasso for T and $xy\in\mathscr{L}$.*
(ii) *$\mathscr{L}$ is an equidistant lasso for T if and only if $\mathscr{L}_1(x,y)\cup\{xy\}$ is an equidistant lasso for T and $xy\in\mathscr{L}$.*
(ii) *If $\mathscr{L}\neq\emptyset$ then $\mathscr{L}$ is a weak lasso for T if and only if $\mathscr{L}_1(x,y)\cup\{xy\}$ is a weak lasso for T and $xy\in\mathscr{L}$.*

Proof. (i): Put $\mathscr{L}_1^+ := \mathscr{L}_1(x,y)\cup\{xy\}$ and assume first that $\mathscr{L}$ is a topological lasso for T. Then $\mathscr{L}\neq\emptyset$. Let T' denote an X-tree and ω and ω' equidistant, proper edge-weightings for T and T', respectively, so that (T,ω) and (T',ω') are $\mathscr{L}_1^+$-isometric. To see that T and T' are equivalent it clearly suffices to show that (T,ω) and (T',ω') are also $\mathscr{L}$-isometric, that is,

$$D_{(T,\omega)}(a,b) = D_{(T',\omega')}(a,b) \tag{2}$$

holds for all $ab\in\mathscr{L}$.

Suppose $ab\in\mathscr{L}$. If $x\notin\{a,b\}$ then $ab\in\mathscr{L}_1^+$ and so Equation (2) holds as (T,ω) and (T',ω') are $\mathscr{L}_1^+$-isometric. So assume that $x\in\{a,b\}$, say $x=a$. If $y=b$ then $ab=xy\in\mathscr{L}$ and so Equation (2) holds by the same argument. If $y\neq b$ then $xb=ab\in\mathscr{L}$ and so $yb\in\mathscr{L}_1^+$. Since $xy\in\mathscr{L}_1^+$ also holds, we have $D_{(T,\omega)}(x,b)=D_{(T',\omega')}(x,b)$ by Lemma 3.1 and so Equation (2) follows in this case, too.

Conversely, suppose that $\mathscr{L}_1^+$ is a topological lasso for T and that $xy\in\mathscr{L}$. Then $\mathscr{L}_1^+\neq\emptyset$. Assume that T' is an X-tree and that ω and ω' are equidistant, proper edge-weightings for T and T', respectively, so that (T,ω) and (T',ω') are $\mathscr{L}$-isometric. To see that T and T' are equivalent it clearly suffices to show that (T,ω) and (T',ω') are also $\mathscr{L}_1^+$-isometric, that is, that

$$D_{(T,\omega)}(a,b) = D_{(T',\omega')}(a,b) \tag{3}$$

holds for all $ab\in\mathscr{L}_1^+$.

Suppose $ab\in\mathscr{L}_1^+$. If $x\in\{a,b\}$ then $ab=xy\in\mathscr{L}$ and so Equation (3) holds as (T,ω) and (T',ω') are $\mathscr{L}$-isometric. So assume that $x\notin\{a,b\}$. Then $ab\in\mathscr{L}_1(x,y)$. If $y\notin\{a,b\}$ holds too then $ab\in\mathscr{L}$ and so Equation (3) holds by the same argument. So assume that $y\in\{a,b\}$, say, $y=a$. Then $yb=ab\in\mathscr{L}_1^+$ and so one of $yb\in\mathscr{L}$ and $xb\in\mathscr{L}$ must hold by the definition of $\mathscr{L}_1^+$. If the former holds then $ab=yb\in\mathscr{L}$ and so Equation (3) holds by assumption on (T,ω) and (T',ω'). If $xb\in\mathscr{L}$ then $D_{(T,\omega)}(y,b)=D_{(T',\omega')}(y,b)$ follows by Lemma 3.1 since, by assumption, $xy\in\mathscr{L}$. But then Equation (3) holds in this case, too.

(ii) & (iii): This follows using similar arguments as in the proof of (i). □

4. The child-edge graph

In this section we first introduce the child-edge graph $G_T(\mathscr{L},v)$ associated to an interior vertex v of an X-tree T and a non-empty set $\mathscr{L}\subseteq\binom{X}{2}$ of cords and then study some of its properties with regards to coralling an X-tree. We start with a definition.

Suppose T is an X-tree, $v\in V^o(T)$, and $\mathscr{L}\subseteq\binom{X}{2}$ is a non-empty set of cords. Then we call the graph $G_T(\mathscr{L},v)=(V_{T,v},E_{T,v})$ with vertex set $V_{T,v}$ the set of all child edges of v and edge set $E_{T,v}$ the set of all $\{e,e'\}\in\binom{V_{T,v}}{2}$ for which there exist leaves $a,b\in X$ such that e and e' are edges on the path from a to b in T and $ab\in\mathscr{L}$ the *child-edge graph (of v with respect to T and $\mathscr{L}$)*. Note that in case there is no danger of ambiguity with regards to the X-tree T we are referring to, we will write $G(\mathscr{L},v)$ rather than $G_T(\mathscr{L},v)$ and V_v and E_v rather than $V_{T,v}$ and $E_{T,v}$.

For T an X-tree, the next result provides a key insight into the structure of $G(\mathscr{L},v)$, $v \in V^o(T)$, whenever $\mathscr{L}$ coralls T. To state it, we denote by $V^o(T)^-$ the set of all interior vertices of T that are not a parent of a pseudo-cherry of T.

Lemma 4.1. *Suppose that T is an X-tree but not the star-tree on X, that $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords that coralls T, and that $v \in V^o(T)$. Then the following holds.*

(i) *If $v \in V^o(T)^-$ and $e \in V_v$ is a child edge of v that is not incident with a leaf of T then $\{e,e'\} \in E_v$, for all $e' \in V_v$ that are incident with a leaf of T. In particular, $G(\mathscr{L},v)$ is connected.*

(ii) *If $v \in V^o(T) - V^o(T)^-$ then $G(\mathscr{L},v)$ is connected.*

Proof. (i): Let $e \in V_v$ denote a child edge of v that is not incident with a leaf of T and let $u \in V(T)$ denote the child of v that is incident with e. Let $\omega : E(T) \to \mathbb{R}_{\geq 0}$ be an equidistant, proper edge-weighting for T. Assume for contradiction that there exists a child $u' \in V(T)$ of v that is a leaf such that with $e_{u'}$ denoting the child edge of v incident with u', we have that $\{e,e_{u'}\} \notin E_v$. Note that since ω is equidistant and u' is a leaf of T whereas u is not, we must have $\omega(e_{u'}) > \omega(e)$.

Assume first that $|V_v| \geq 3$. Let T' denote the X-tree obtained from T by deleting the edge $e_{u'}$ and adding the edge $e^* = \{u,u'\}$. Clearly, T' is not a refinement of T. Consider the edge-weighting

$$\omega' : E(T') \to \mathbb{R}_{\geq 0} : \ f \mapsto \begin{cases} \omega(f) & \text{if } f \neq e^*, \\ \omega(e_{u'}) - \omega(e) & \text{else.} \end{cases}$$

Then it is easy to see that ω' is equidistant and proper. Since, by construction, (T,ω) and (T',ω') must be $\mathscr{L}$-isometric it follows that T' is a refinement of T as $\mathscr{L}$ coralls T; a contradiction.

Now assume that $|V_v| = 2$. If $v \neq \rho_T$ then v must have a parent $w \in V(T)$. Let T' denote the X-tree obtained from T as before except that we now suppress v as this has rendered it a vertex with a single child. Let ω' be the edge-weighting for T' as defined above except that we put $\omega'(\{u,w\}) = \omega(\{u,v\}) + \omega(\{v,w\})$. Then the same arguments as in the previous case yield a contradiction

If $v = \rho_T$ then let T' denote the X-tree obtained from T by collapsing the edge $\{v,u\}$. Clearly, T' is not a refinement of T. Consider the edge-weighting $\omega' : E(T') \to \mathbb{R}_{\geq 0}$ for T' defined by putting $\omega' = \omega|_{E(T')}$. Then the same arguments as in the previous two cases yield a contradiction.

(ii): Assume for contradiction that there exists some $v \in V^o(T) - V^o(T)^-$ such that $G(\mathscr{L},v)$ is not connected. Then every child of v is a leaf of T and there exist vertices $e_1,e_2 \in V_v$ distinct such that e_1 and e_2 are not joined by a path in $G(\mathscr{L},v)$. Let G_1 and G_2 denote the connected components of $G(\mathscr{L},v)$ containing e_1 and e_2, respectively. For all children $u \in V(T)$ of v, let e_u denote the child edge of v incident with u. Note that since T is not the star-tree on X there must exist a vertex $w \in V(T)$ that is the parent of v. Let T' denote the X-tree obtained from T via the following process. Let $i = 1,2$. Then, for all $u_i \in V(G_i)$, subdivide the edge e_{u_i} by a new vertex p_{u_i} not already contained in $V(T)$. Next, identify all vertices p_{u_i} into the vertex p_i and then delete all copies of the edges $\{v,p_i\}$ from T. Finally, add the edge $\{w,p_i\}$ and suppress p_i if $|V(G_i)| = 1$. If $V_v = V(G_1) \cup V(G_2)$ then also suppress the vertex v. The resulting tree is T' and, in either case, T' is clearly not a refinement of T.

Let ω denote an equidistant, proper edge-weighting for T. Note that $\omega(e_1) = \omega(e)$ must hold for all $e \in V_v$, as $v \in V^o(T) - V^o(T)^-$. For the following, assume first that neither p_1

nor p_2 have been suppressed in the construction of T'. Consider the edge-weighting

$$\omega' : E(T') \to \mathbb{R}_{\geq 0} : \ e \mapsto \begin{cases} \omega(e) & \text{if } p_1, p_2 \notin e, \\ \omega(\{v,w\}) & \text{if } e \in \{\{w,p_1\},\{w,p_2\}\} \\ \omega(e_1) & \text{else.} \end{cases}$$

Then, by construction, ω' is equidistant and proper and (T,ω) and $T',\omega')$ are $\mathscr{L}$-isometric. Since, by assumption, $\mathscr{L}$ coralls T it follows that T' is a refinement of T; a contradiction.

In case one of p_1 and p_2 or both of them have been suppressed in the construction of T' the definition of the edge-weighting ω' for T' is similar to the one above thus leading to a contradiction in these cases too. □

The next result is a strengthening of Lemma 4.1(i). To state it, we require further terminology concerning child-edge graphs. Suppose T is an X-tree, $v \in V^o(T)^-$, and $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords. Then we denote by $E_l(v) \subseteq E(T)$ the set of child edges of v that are incident with a leaf of T and by $E_s(v) \subseteq E(T)$ the set of child edges of v that are not contained in $E_l(v)$. Note that $E_s(v)$ is empty if and only if v is the parent of a pseudo-cherry of T. Also note that $E_l(v) = \emptyset$ might hold. Clearly, if neither of them is the empty-set then $\{E_l(v), E_s(v)\}$ is a partition of V_v. For $v \in V^o(T)^-$, we say that $G(\mathscr{L},v)$ is *rich* if the subgraph $G_T(\mathscr{L},v)_s$ of $G_T(\mathscr{L},v)$ induced by $E_s(v)$ is a clique, and, in case $E_l(v) \neq \emptyset$, we have for all $e \in E_l(v)$ and all $e' \in E_s(v)$ that $\{e,e'\} \in E_v$. As before we will write $G(\mathscr{L},v)_s$ rather than $G_T(\mathscr{L},v)_s$ if there is no ambiguity with regards to which X-tee T we are referring. Note that with T and v as above, if $E_l(v) = \emptyset$ then $G(\mathscr{L},v)$ is rich if and only if $G(\mathscr{L},v)_s$ is a clique.

Proposition 4.2. *Suppose that T is an X-tree but not the star-tree and that $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords. If $\mathscr{L}$ coralls T then $G(\mathscr{L},v)$ must be rich, for all vertices $v \in V^o(T)^-$.*

Proof. Let ω denote an equidistant, proper edge-weighting for T and assume for contradiction that there exists a vertex $v \in V^o(T)^-$ such that $G(\mathscr{L},v)$ is not rich. We first show that $G(\mathscr{L},v)_s$ must be a clique. Suppose $G(\mathscr{L},v)_s$ is not a clique, that is, there exist distinct child edges e and e' of v contained in $V(G(\mathscr{L},v)_s)$ such that with v_e and $v_{e'}$ denoting the children of v incident with e and e', respectively, we have, for all $a \in L(v_e)$ and all $b \in L(v_{e'})$ that $ab \notin \mathscr{L}$. Note that $|V_v| \geq 3$ must hold. Indeed, if $|V_v| = 2$ then $V_v = \{e,e'\}$ and so $\{e,e'\} \in E_v$ since, by Lemma 4.1, $G(\mathscr{L},v)$ is connected. But then $G(\mathscr{L},v)$ is a clique and so $G(\mathscr{L},v)_s$ is a clique; a contradiction. Without loss of generality assume that $\omega(e) \geq \omega(e')$.

If $\omega(e) > \omega(e')$ then consider the X-tree T' obtained from T by deleting the edge $\{v,v_e\}$ and attaching v_e to $v_{e'}$ via the edge $e^* = \{v_e, v_{e'}\}$. Clearly, T' is not a refinement of T. Consider the edge-weighting ω' for T' defined by putting

$$\omega' : E(T') \to \mathbb{R}_{\geq 0} : \ f \mapsto \begin{cases} \omega(f) & \text{if } f \neq e^*, \\ \omega(e) - \omega(e') & \text{else.} \end{cases}$$

Clearly, ω' is equidistant and proper and, by construction, (T,ω) and (T',ω') are $\mathscr{L}$-isometric. Since $\mathscr{L}$ coralls T it follows that T' must be a refinement; a contradiction.

If $\omega(e) = \omega(e')$ then consider the X-tree T' obtained from T by first identifying the vertices v_e and $v_{e'}$ (keeping the label v_e) and then deleting one of the edges from v to v_e. Again, T' is clearly not a refinement of T. Consider the edge-weighting $\omega' : E(T') \to \mathbb{R}_{\geq 0}$ defined as $\omega' = \omega|_{E(T')}$. Then the same arguments as before imply that T' is a refinement of T; a contradiction. Thus, $G(\mathscr{L},v)_s$ is a clique, as required.

Now if $E_l(v) = \emptyset$ then $G(\mathscr{L},v)$ must be rich. So assume that $E_l(v) \neq \emptyset$. But then Lemma 4.1(i), implies that $G(\mathscr{L},v)$ must be rich. □

We conclude this section with a result that will be useful for establishing the aforementioned characterization of weak lassos in terms of child-edge graphs (Theorem 5.1). Its proof relies on the well-known fact that an X-tee T' is a refinement of an X-tree T if and only if $\mathscr{R}(T) \subseteq \mathscr{R}(T')$ (see e.g. (Semple and Steel, 2003, Theorem 6.4.1)).

Lemma 4.3. *Suppose that T is an X-tree that has a unique cherry x,y and that $\mathscr{L} \subseteq \binom{X}{2}$ is a set of cords that contains the set $\{xy\} \cup \{az : z \in X - \{x,y\}$ and $a = x$ or $a = y\}$. Then $\mathscr{L}$ coralls T.*

Proof. By Theorem 3.2, it suffices to show that $\mathscr{L}' := \mathscr{L}(x,y) \cup \{xy\}$ coralls T. Suppose there exists an X-tree T' and equidistant, proper edge-weightings ω and ω' of T and T', respectively, such that (T,ω) and (T',ω') are $\mathscr{L}'$-isometric. To see that T' is a refinement of T it suffices to show that $xy|z \in \mathscr{R}(T')$ holds for all $z \in X - \{x,y\}$. Let $z \in X - \{x,y\}$. Then $zy \in \mathscr{L}'$. Combined with the facts that $xy \in \mathscr{L}'$ and that $xy|z \in \mathscr{R}(T)$ it follows, by Lemma 3.1, that $xy|z \in \mathscr{R}(T')$, as required. □

5. A CHARACTERIZATION OF A WEAK LASSO

In this section, we characterize sets of cords of X that corall an X-tree T in terms of two properties on the child edge-graphs associated to the interior vertex of T. In addition, we present a simple example that illustrates that this characterization does not hold if the equidistant requirement for the two edge-weight functions mentioned in the definition of such a lasso is dropped.

Theorem 5.1. *Suppose that T is an X-tree but not the star-tree and that $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords. Then $\mathscr{L}$ is a weak lasso for T if and only if the following two properties hold*

(C1) *$G(\mathscr{L},v)$ is rich, for all $v \in V^o(T)^-$*
(C2) *$G(\mathscr{L},v)$ is connected, for all $v \in V^o(T) - V^o(T)^-$.*

Proof. Assume first that $\mathscr{L}$ coralls T. Then Properties (C1) and (C2) must hold by Proposition 4.2 and Lemma 4.1(ii).

To see the converse, assume that Properties (C1) and (C2) hold. We prove that $\mathscr{L}$ must be a weak lasso for T by induction on the size n of X. Note that the statement clearly holds in case $n = 3$ as then T is binary and a refinement of a binary X-tree is the tree itself. So assume that $n = 4$. Then since T is not the star-tree on X, we have $2 \leq |V^o(T)| \leq 3$. If $|V^o(T)| = 3$ then T is binary and so the statement holds. So assume that $|V^o(T)| = 2$. Then T has a unique cherry x,y and so, by Lemma 4.3, $\mathscr{L}$ coralls T in this case too.

Assume that the statement holds for all finite sets of size $n \geq 4$ and let X denote a set of size $n+1$. Let T be an X-tree that is not the star-tree on X and let $\mathscr{L} \subseteq \binom{X}{2}$ denote a non-empty set of cords such that Properties (C1) and (C2) are satisfied for T and $\mathscr{L}$. Note that T must contain at least one pseudo-cherry. To see that $\mathscr{L}$ coralls T, let T' denote an X-tree and ω and ω' equidistant, proper edge-weightings for T and T', respectively, such that (T,ω) and (T',ω') are $\mathscr{L}$-isometric. We distinguish between the cases that (i) every pseudo-cherry of T is in fact a cherry of T and (ii) that T contains a pseudo-cherry that has at least three leaves.

Case (i): Assume that every pseudo-cherry of T is a cherry and let $x,y \in X$ such that x,y is a cherry of T. Note that since $n \geq 5$ and T is not the star-tree on X, there must exist a vertex $w \in V(T)$ that is the parent of v_x (which is itself the parent of the cherry x,y). Put

$X_1 = X - \{x\}$ and $\mathscr{L}_1 = \mathscr{L}_1(x,y)$. Note that since $G(\mathscr{L}, v_x)$ is connected, it immediately follows that $xy \in \mathscr{L}$ and that since $G(\mathscr{L}, w)$ is rich we must have $\mathscr{L}_1 \neq \emptyset$. Let T_1 denote the X_1-tree obtained from T by deleting x and its incident edge and suppressing v_x as this has rendered it a vertex with a single child. Let

$$\omega_1 : E(T_1) \to \mathbb{R}_{\geq 0} : e \mapsto \begin{cases} \omega(e) & \text{if } e \neq \{w,y\} \\ \omega(\{w,v_x\}) + \omega(\{v_x,y\}) & \text{else.} \end{cases}$$

Clearly, ω_1 is equidistant and proper and T_1 is either the star-tree on X_1 or not. Assume first that T_1 is the star-tree on X_1. Then x,y is the unique cherry of T and all elements $z \in X_1 - \{y\}$ are adjacent with the root ρ_T of T which is w. But then Properties (C1) and (C2) combined with Lemma 4.3 imply that $\mathscr{L}$ coralls T.

So assume that T_1 is not the star-tree on X_1. We claim that Properties (C1) and (C2) hold for T_1 and $\mathscr{L}_1$. We start with establishing Property (C1). Assume for contradiction that there exists some $u \in V^o(T_1)^-$ such that $G_{T_1}(\mathscr{L}_1, u)$ is not rich. We first show that $G_{T_1}(\mathscr{L}_1, u)_s$ must be a clique. Assume for contradiction that $G_{T_1}(\mathscr{L}_1, u)_s$ is not a clique, that is, there exist vertices e and e' in $G_{T_1}(\mathscr{L}_1, u)_s$ such that for all $a \in L_{T_1}(u_e)$ and all $b \in L_{T_1}(u_{e'})$ we have that $ab \notin \mathscr{L}_1$ where u_e and $u_{e'}$ denote the children of u in T_1 incident with e and e', respectively. Note that $V^o(T_1) \cup \{v_x\} = V^o(T)$ and that $u = w$ must hold. Indeed assume for contradiction that $u \neq w$. Then v_x is not a child of u in T and so u_e and $u_{e'}$ are also children of u in T. Since $G_T(\mathscr{L}, u)_s$ is a clique by Property (C1), there must exist $a \in L_T(u_e)$ and $b \in L_T(u_{e'})$ such that $ab \in \mathscr{L}$. If $x \notin \{a,b\}$ then, by the definition of $\mathscr{L}_1$, we have $ab \in \mathscr{L}_1$; a contradiction. Thus, $x \in \{a,b\}$. Without loss of generality assume that $x = a$. Then $y \in L_T(u_e)$ and, again by the definition of $\mathscr{L}_1$, we obtain $yb \in \mathscr{L}_1$; a contradiction. Thus $u = w$, as required. But then $y \in \{u_e, u_{e'}\}$ and so one of e and e' is not a vertex in $G_{T_1}(\mathscr{L}_1, u)_s$; a contradiction. Thus, $G_{T_1}(\mathscr{L}_1, u)_s$ must be a clique, as required.

Since, by assumption, $G_{T_1}(\mathscr{L}_1, u)$ is not rich, there must therefore exist a leaf z of T_1 with $e' = \{u,z\} \in E(T_1)$ holding and some vertex e in $G_{T_1}(\mathscr{L}_1, u)_s$ such that $\{e,e'\}$ is not an edge in $G_{T_1}(\mathscr{L}_1, u)$. Let u_e denote the child of u in T_1 incident with e. If $u \neq w$ then since the children of u in T are precisely the children of u in T_1 and, by Property (C1), $G_T(\mathscr{L}, u)$ is rich we obtain a contradiction. Thus, $u = w$. But then $y = z$ must hold. Since $G_T(\mathscr{L}, w)$ is rich there must exist some $a \in \{x,y\}$ and some $b \in L(u_e)$ such that $ab \in \mathscr{L}$. But then $yb \in \mathscr{L}_1$ and so $\{e,e'\}$ is an edge in $G_{T_1}(\mathscr{L}_1, u)$, a contradiction.

We next establish that Property (C2) is satisfied by T_1 and $\mathscr{L}_1$ which will conclude the proof of the claim. Let $u \in V^o(T_1) - V^o(T_1)^-$. Then u must be the parent of a pseudo-cherry of T_1. If $u \neq w$ then since, by assumption, every pseudo-cherry of T is a cherry of T it follows that u is the parent of a cherry of T_1. But then $G_{T_1}(\mathscr{L}_1, u)$ is connected as $G_{T_1}(\mathscr{L}_1, u) = G_T(\mathscr{L}, u)$ and Property (C2) is satisfied by T and $\mathscr{L}$. So assume that $u = w$. Then u is the parent of v_x in T and all children of u in T but v_x are leaves of T. Since, by Property (C1), $G_T(\mathscr{L}, u)$ is rich, there exists for all children z of u that are leaves of T some $b_z \in \{x,y\}$ such that $b_z z \in \mathscr{L}$. But then $yz \in \mathscr{L}_1$ for all such children z of u and thus $G_{T_1}(\mathscr{L}_1, u)$ is connected, as required. Thus Property (C2) is also satisfied by T_1 and $\mathscr{L}_1$ which completes the proof of the claim. By induction, it follows that $\mathscr{L}_1$ is a weak lasso for T_1.

Let T_1' denote the X_1-tree obtained from T' by deleting x and its incident edge and suppressing the parent vertex of x in T_1' if this has rendered it a vertex with a single child. Let ω_1' denote the edge-weighting of T_1' that is canonically induced by ω_1 on the edges of T_1'. Then, by Lemma 3.1 combined with the assumption that (T, ω) and (T', ω') are

$\mathscr{L}$-isometric, it follows that (T_1,ω_1) and (T_1',ω_1') are $\mathscr{L}_1$-isometric. Since $\mathscr{L}_1$ is a weak lasso for T_1 this implies that T_1' is a refinement for T_1.

To establish that $\mathscr{L}$ coralls T it now suffices to show that $xy|a \in \mathscr{R}(T')$ holds for all $a \in L(w) - \{x,y\}$. To this end, note that since Property (C1) is satisfied by T and $\mathscr{L}$, we have for all children $a \in L(w)$ that are leaves of T that there exists some $b \in L_T(v_x) = \{x,y\}$ such that $ab \in \mathscr{L}$. Combined with Lemma 3.1 and $xy \in \mathscr{L}$, it follows that $D_{(T,\omega)}(x,a) = D_{(T',\omega')}(x,a) = D_{(T,\omega)}(y,a) = D_{(T',\omega')}(y,a)$. Since $xy|a \in \mathscr{R}(T)$ we obtain $xy|a \in \mathscr{R}(T')$, as required. This completes the proof of the induction step in this case.

Case (ii): Assume that T contains a pseudo-cherry $\mathfrak{c}$ of size three or more. Let $v \in V^o(T)$ denote the parent of $\mathfrak{c}$ in T. Then, by Property (C2), $G(\mathscr{L},v)$ is connected. By (Diestel, Electronic Edition 2005, Proposition 1.4.1), we may choose some $x \in X$ such that the graph $G^-(\mathscr{L},v)$ obtained from $G(\mathscr{L},v)$ by deleting e_x and its incident edges is connected. Note that since x is a leaf in $\mathfrak{c}$, we have $v = v_x$.

Put $X_1 := X - \{x\}$, choose some $y \in L(v_x)$ such that $xy \in \mathscr{L}$, and put $\mathscr{L}_1 := \mathscr{L}(x,y)$. Consider the X-tree T_1 obtained from T by deleting x and its incident edge. We claim again that $G_{T_1}(\mathscr{L}_1,u)$ satisfies Properties (C1) and (C2) for all $u \in V^o(T_1)$ as specified in those conditions. To see this, note first that since v_x has at least two children in T_1, we have $V^o(T_1) = V^o(T)$ and so also $V^o(T_1)^- = V^o(T)^-$.

We start with establishing Property (C1). Assume for contradiction that there exists some vertex $u \in V^o(T_1)^-$ such that $G_{T_1}(\mathscr{L}_1,u)$ is not rich. We again show first that $G_{T_1}(\mathscr{L}_1,u)_s$ is a clique. Assume for contradiction that this is not the case and let e, e', u_e, and $u_{e'}$ be as in the corresponding situation in the previous case. Note that since $u \in V^o(T_1)^- = V^o(T)^-$ and $v_x \in V^o(T) - V^o(T)^-$ we have $v_x \neq u$. But then similar arguments as the ones used to show in Case (i) that $w \neq u$ in the context of establishing that $G_{T_1}(\mathscr{L}_1,u)_s$ is a clique yield a contradiction. Thus, $G_{T_1}(\mathscr{L}_1,u)_s$ must be a clique, as required. As in the previous case, there must therefore exist some $z \in X_1$ such that $e = \{z,u\} \in E(T_1)$ and some vertex e' in $G_{T_1}(\mathscr{L}_1,u)_s$ such that $\{e,e'\}$ is not an edge in $G_{T_1}(\mathscr{L}_1,u)$. Note that the same arguments as above imply that $u \neq v_x$. By the definition of $\mathscr{L}_1$, it follows that x must be the unique leaf in $L_T(u_{e'})$ such that $xz \in \mathscr{L}$. Since y and x are leaves in the same pseudo-cherry of T, we obtain $yz \in \mathscr{L}_1$. Consequently, $\{e,e'\}$ is an edge in $G_{T_1}(\mathscr{L}_1,u)$, a contradiction. Thus, Property (C1) is satisfied by T_1 and $\mathscr{L}_1$.

To see that T_1 and $\mathscr{L}_1$ satisfy Property (C2) assume without loss of generality that there exists some $u \in V^o(T) - V^o(T)^-$ such that $G_{T_1}(\mathscr{L}_1,u)$ is not connected. Then $u \neq v_x$, by the choice of x. Since u is the parent of a pseudo-cherry in T it follows that u must be the parent of the same pseudo-cherry in T_1. But then $G_{T_1}(\mathscr{L}_1,u) = G_T(\mathscr{L},u)$ and so $G_{T_1}(\mathscr{L}_1,u)$ must be connected as, by Property (C2), $G_T(\mathscr{L},u)$ is connected; a contradiction. This concludes the proof of the claim. By induction, it follows that $\mathscr{L}_1$ coralls T_1.

Let T_1', w, and ω_1' be as in the previous case. Then, as in that case, T_1' must be a refinement of T_1. We claim that for all children u of w in T distinct from v_x and all $a \in L_T(u)$ we must have

$$xy|a \in \mathscr{R}(T').$$

To see this, note first that Property (C1) implies for all such children u of w that there must exist some $a_u \in L_T(u)$ and some $z_{v_x} \in L_T(v_x)$ such that $a_u z_{v_x} \in \mathscr{L}$. Let u denote a child of w in T distinct from v_x and put $a = a_u$ and $z = z_{v_x}$. We show first that

$$D_{(T,\omega)}(a,y) = D_{(T',\omega')}(a,y) \text{ and } D_{(T,\omega)}(a,x) = D_{(T',\omega')}(a,x). \tag{4}$$

Clearly, if $z \in \{x,y\}$ then one of the two equations in (4) must hold. Assume without loss of generality that $z \neq x$. Since $G_T(\mathscr{L},v_x)$ is connected by Property (C2), it follows that there

exists a path $z = z_1, z_2, \ldots, z_k = x$, $k \geq 2$, from z to x in $G_T(\mathscr{L}, v_x)$. But then Lemma 3.1(i) implies $D_{(T,\omega)}(a,x) = D_{(T',\omega')}(a,x)$ since, for all $1 \leq i \leq k-1$, we have $z_i z_{i+1} \in \mathscr{L}$. If $z \neq y$ then similar arguments imply that $D_{(T,\omega)}(a,y) = D_{(T',\omega')}(a,y)$. If $z = y$ then $ya = za \in \mathscr{L}$ and so $D_{(T,\omega)}(a,y) = D_{(T',\omega')}(a,y)$. Thus both equations in (4) must hold, as required. Combined with $xy \in \mathscr{L}$ (which holds by the choice of y) and the fact that $xy|a \in \mathscr{R}(T)$ we obtain

$$xy|a \in \mathscr{R}(T')$$

in view of Lemma 3.1(ii). Thus the claim follows if $|L_T(u)| = 1$. So assume that $|L_T(u)| \geq 2$. Suppose $a' \in L_T(u) - \{a\}$. Then $y|a'a \in \mathscr{R}(T_1')$ must hold as T_1' is a refinement of T_1 and $y|a'a \in \mathscr{R}(T_1)$. Since the only $\{x,y,a',a\}$-tree that can simultaneously display the triplets $y|a'a$ and $xy|a$ is the tree with cherries x,y and a',a and that tree is equivalent with the tree $T|_{\{x,y,a',a\}}$ it follows that $xy|a' \in \mathscr{R}(T')$, as claimed.

Combined with the fact that T_1' is a refinement of T_1 it follows that T' is a refinement of T. Hence, $\mathscr{L}$ coralls T which concludes the proof of the induction step in this case too and, thus, the proof of the theorem.

□

Note that Theorem 5.1 immediately implies that for a non-empty set $\mathscr{L}$ of cords of X to be a weak lasso for an X-tree T that is not the star-tree on X, it must be a *covering* of X, that is, $X = \bigcup_{A \in \mathscr{L}} A$. Also note that it immediately implies that a minimum size weak lasso for T must have $\sum_{v \in V^o(T)^-} (\binom{|V(G(\mathscr{L},v)_s)|}{2} + |V(G(\mathscr{L},v)_s)| \times |V(G(\mathscr{L},v)_l)|) + \sum_{v \in V^o(T) - V^o(T)^-} |V_v|$ cords. Thus, such a lasso has at most $|V^o(T)^-|\binom{m}{2} + (m-1)|V^o(T) - V^o(T)^-|$ cords where $m = \max_{v \in V^o(T)^-} |V_v|$ and at least $(l-1)|V^o(T)|$ cords where $l = \min_{v \in V^o(T)^-} |V_v|$. Note that these bounds are sharp in the case that all interior vertices of T have the same number k of children. In the former case T is such that no interior vertex of T that is not a parent of a pseudo-cherry is adjacent with a leaf of T. In the latter case T is the *bearded caterpillar tree* on X, that is, T is a (rooted) path and every vertex of that path is adjacent with $k-1$ leaves but the end vertex of T that is not ρ_T which has k children.

Finally, note that as the example presented in Fig. 2 illustrates, for the theorem to hold the requirement that both proper edge-weightings are equidistant in the definition of a weak lasso cannot be dropped. More precisely for $X = \{a,b,c,d,e,f\}$ and $\mathscr{L} = \{ab, bc, bd, af, ae\}$ the X-tree T pictured on the left of that figure (with the indicated edge-weighting ω ignored for the moment) satisfies Properties (C1) and (C2) and the X-tree T' depicted on the right of that figure (again with the indicated edge-weighting ω' ignored where $0 < \varepsilon < 1$) is clearly not a refinement of T. Also ω and ω' are obviously proper and, in the case of ω, equidistant and (T,ω) and (T',ω') are $\mathscr{L}$-isometric.

6. A CHARACTERIZATION OF AN EQUIDISTANT LASSO

In this section, we present a characterization of an equidistant lasso $\mathscr{L} \subseteq \binom{X}{2}$ for an X-tree T in terms of the child-edge graphs associated to the interior vertices of T. To establish it, we require a further notation. Suppose T is an X-tree and v and w are two vertices of T. Then we denote by $E_T(v,w)$ the set of all edges of T on the path from v to w.

Theorem 6.1. *Suppose T is an X-tree and $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords. Then the following are equivalent:*

(i) *$\mathscr{L}$ is an equidistant lasso for T.*
(ii) *for every vertex $v \in V^o(T)$, the graph $G(\mathscr{L},v)$ contains at least one edge.*




FIGURE 2. An example illustrating that for Theorem 5.1 to hold the equidistance requirement in the definition of a weak lasso cannot be dropped (see text for details).

Proof. (i) $\Rightarrow$ (ii): Suppose $\mathscr{L}$ is an equidistant lasso for T and assume for contradiction that there exists an interior vertex v of T for which $G(\mathscr{L},v)$ does not contain an edge. Note first that $v = \rho_T$ cannot hold. Indeed, suppose ω_1 is a equidistant, proper edge-weighting for T and let $\varepsilon > 0$ be a sufficiently small real number. Consider the edge-weighting $\omega_2 : E(T) \to \mathbb{R}_{\geq 0}$ defined by putting, for all $e \in E(T)$, $\omega_2(e) = \omega_1(e) - \varepsilon$ if $\rho_T \in e$ and $\omega_2(e) = \omega_1(e)$ else. Clearly, ω_2 is an equidistant, proper edge-weighting for T distinct from ω_1 and (T,ω_1) and (T,ω_2) are $\mathscr{L}$-isometric. Since, by assumption, $\mathscr{L}$ is an equidistant lasso for T it follows that $\omega_1 = \omega_2$; a contradiction. Thus, $v \neq \rho_T$ and so there must exist a parent $w \in V(T)$ of v in T. Let ω_1 denote again an equidistant, proper edge-weighting for T. Consider the map $\omega_2 : E(T) \to \mathbb{R}_{\geq 0}$ defined by putting

$$\omega_2(e) = \begin{cases} \omega_1(e) & \text{if } v \notin e, \\ \omega_1(e) - \varepsilon & \text{if } e = \{v,w\}, \\ \omega_1(e) + \varepsilon & \text{else,} \end{cases}$$

where $\varepsilon > 0$ is small enough. Clearly, ω_2 is an equidistant, proper edge-weighting for T which is distinct from ω_1. By construction, (T,ω_1) and (T,ω_2) are $\mathscr{L}$-isometric and so $\omega_1 = \omega_2$ must hold as $\mathscr{L}$ is an equidistant lasso for T; a contradiction.

(ii) $\Rightarrow$ (i): Suppose that, for all $v \in V^o(T)$, the graph $G(\mathscr{L},v)$ has at least one edge and assume for contradiction that $\mathscr{L}$ is not an equidistant lasso for T. Then there exist distinct equidistant, proper edge-weightings ω_1 and ω_2 for T such (T,ω_1) and (T,ω_2) are $\mathscr{L}$-isometric. Thus, there must exist some edge $e \in E(T)$ such that $\omega_1(e) \neq \omega_2(e)$.

Assume without loss of generality that $e = \{v, v_0\}$ is such that with v being the parent of v_0 we have that $\omega_1(e') = \omega_2(e')$ holds for all edges e' of T that lie on a path from v_0 to a leaf of T contained in $L(v_0)$. Note that v_0 could be a leaf of T in which case such a path has length zero. Let $v_1, v_2, \ldots, v_l$, $l \geq 1$ denote the other children of v. Then, by assumption, $G(\mathscr{L},v)$ contains at least one edge and so there exist $i,j \in \{0,l\}$ distinct and leaves $y_i, y_j \in L(v)$ that are descendants of v_i and v_j (or coincide with them), respectively, such that $y_i y_j \in \mathscr{L}$. Let $z \in L(v_0)$ denote a leaf of T. By Property (E1), we obtain

$$\begin{aligned} \omega_1(e) + \sum_{e' \in E_T(v_0,z)} \omega_1(e') &= D_{(T,\omega_1)}(v,z) = \frac{1}{2} D_{(T,\omega_1)}(y_i,y_j) = \frac{1}{2} D_{(T,\omega_2)}(y_i,y_j) \\ &= D_{(T,\omega_2)}(v,z) = \sum_{e' \in E_T(v_0,z)} \omega_2(e') + \omega_2(e). \end{aligned}$$

and so $\omega_1(e) = \omega_2(e)$ follows by the choice of e; a contradiction. Thus, $\mathscr{L}$ must be an equidistant lasso for T. $\square$

Theorem 6.1 immediately implies that an equidistant lasso on X need not be a covering of X. Also, it implies that the size of a minimum equidistant lasso for an X-tree T is $|V^\circ(T)|$. For the extreme cases that T is the star tree on X such a lasso has precisely one element and if T is binary such a lasso has $|X|-1$ elements as any such tree is known to have $|X|-1$ interior vertices (see e. g. Semple and Steel (2003)).

Theorem 5.1 combined with Theorem 6.1 immediately implies the following link between equidistant and weak lassos.

Corollary 6.2. *Suppose T is an X-tree and $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords. Then $\mathscr{L}$ is an equidistant lasso for T whenever it is a weak lasso for T.*

We remark in passing that dropping the requirement that the two edge-weightings have to be equidistant in the definition of an equidistant lasso gives rise to the definition of an *edge-weight lasso* for an X-tree. However it is easy to see that Theorem 6.1 does not hold with equidistant lasso replaced by edge-weight lasso.

7. A CHARACTERIZATION OF A TOPOLOGICAL LASSO

In this section, we prove the companion result for when a set of cords is a topological lasso for an X-tree T in terms of the child-edge graphs associated to the interior vertices of T. We start again with some more notation.

Suppose that T is an X-tree and that $v \in V^o(T)$ but not the root of T. Then we denote the $L(v)$-tree obtained from T by deleting the parent edge of v by $T_{L(v)}$. Now suppose that $Y \subsetneq X$ is such that there exists some $v \in V^o(T)$ such that $Y = L(v)$. Then we denote the root of T_Y by $\rho(T_Y)$.

Theorem 7.1. *Suppose T is an X-tree and $\mathscr{L} \subseteq \binom{X}{2}$ is a non-empty set of cords. Then the following are equivalent:*

(i) *$\mathscr{L}$ is a topological lasso for T.*
(ii) *for every vertex $v \in V^o(T)$, the graph $G(\mathscr{L},v)$ is a clique.*

Proof. (i) $\Rightarrow$ (ii): Suppose that $\mathscr{L}$ is a topological lasso for T and assume for contradiction that there exists some vertex $v \in V^o(T)$ such that $G(\mathscr{L},v)$ is not a clique. Then there exist child edges $e,e' \in E(T)$ of v such that $\{e,e'\} \notin E_v$. Let v_e and $v_{e'}$ denote the children of v incident with e and e', respectively, and let ω denote an equidistant, proper edge-weighting of T. We distinguish the cases that (i) v is the parent of a pseudo-cherry of T and that (ii) $v \in V^o(T)^-$.

Case (i): Assume that v is the parent of a pseudo-cherry of T. Then since every topological lasso for T is in particular a weak lasso for T, Theorem 5.1 implies that $G(\mathscr{L},v)$ is connected and that, in addition to $x := v_e$ and $y := v_{e'}$, there must exists a further child of v (that is a leaf of T). Note that since ω is equidistant, we must have $\omega(e) = \omega(e')$.

Let T' denote the X-tree obtained from T by subdividing the edge e' by a vertex w that is not already contained in $V(T)$, adding the edge $\{x,w\}$ and deleting the edge e. Clearly, T and T' are not equivalent. Let $0 < \varepsilon < \omega(e)$ and consider edge-weighting

$$\omega' : E(T') \to \mathbb{R}_{\geq 0} : \ f \mapsto \begin{cases} \omega(f) & \text{if } w \notin f, \\ \omega(f) - \varepsilon & \text{if } f \in \{\{w,x\},\{w,y\}\}, \\ \varepsilon & \text{else.} \end{cases}$$

Clearly, ω' is equidistant and proper and it is straight forward to see that (T',ω') and (T,ω) are $\mathscr{L}$-isometric. Since $\mathscr{L}$ is a topological lasso for T it follows that T and T' must be equivalent, a contradiction.

Case (ii): Assume that $v \in V^o(T)^-$. Then since every topological lasso for T is in particular a weak lasso for T, Theorem 5.1 implies that $G(\mathscr{L},v)$ is rich. But then v_e and $v_{e'}$ must be leaves of T. With $x = v_e$ and $y = v_{e'}$ we obtain a contradiction using the same arguments as in Case (i).

(ii) $\Rightarrow$ (i): Suppose that, for every vertex $v \in V^o(T)$, the graph $G(\mathscr{L},v)$ is a clique and assume that T' is an X-tree and ω and ω' are equidistant, proper edge-weightings for T and T', respectively, such that (T,ω) and (T,ω') are $\mathscr{L}$-isometric. If T is the star-tree on X then $\mathscr{L}$ must necessarily be a topological lasso for T. So assume that T is not the star-tree on X. Then, by Theorem 5.1, $\mathscr{L}$ is a weak lasso for T and so T' must be a refinement of T.

We next show that T and T' are in fact equivalent. Assume for contradiction that T and T' are not equivalent. Then there must exist a non-empty subset $Y \subsetneq X$ of leaves of T such that the subtree T_Y of T with leaf set Y is equivalent with the subtree T'_Y of T' with leaf set Y but the subtrees of T and T' with root the parents of $\rho(T_Y)$ and $\rho(T'_Y)$, respectively, are not. Let w denote the parent of $\rho(T_Y)$ in T and w' the parent of $\rho(T'_Y)$ in T'. Then there must exist some $z \in L(w') - L(\rho(T'_Y))$ and distinct $x,y \in L(\rho(T'_Y)) = Y$ such that $xy|z \in \mathscr{R}(T')$ and $xy|z \notin \mathscr{R}(T)$. Hence, $\rho(T_Y)$ must lie on the path from x to y in T. Combined with the fact that T' is a refinement of T, it follows that $T|_{\{x,y,z\}}$ is the star-tree on $Z := \{x,y,z\}$ whose unique interior vertex is $\rho(T_Y)$. Since ω is equidistant, we obtain

$$D_{(T,\omega)}(x,y) = D_{(T,\omega)}(x,z) = D_{(T,\omega)}(z,y). \tag{5}$$

Let $e_x, e_y, e_z \in E(T)$ denote the child edges of $\rho(T_Y)$ that are crossed by a path from $\rho(T_Y)$ to x, y, and z, respectively, and let v_{e_s} denote the child of $\rho(T_Y)$ incident with e_s, for all $s \in Z$. Since, by assumption, $G(\mathscr{L},\rho(T_Y))$ is a clique there must exist leaves $a \in L(v_{e_x})$, $b \in L(v_{e_y})$, and $c \in L(v_{e_z})$ such that $ab, bc, ca \in \mathscr{L}$. By the same reason, it follows in view of Lemma 3.1 that $D_{(T,\omega)}(p,q) = D_{(T',\omega')}(p,q)$ must hold for any two elements $p,q \in Z$ distinct. Combined with Equality (5), we obtain $D_{(T',\omega')}(p,q) = D_{(T',\omega')}(p,q)$, for any two such elements p and q. But then $T'|_Z$ must be the star-tree on Z and so $xy|z \notin \mathscr{R}(T')$, a contradiction. □

Note that Theorem 7.1 immediately implies that a topological lasso $\mathscr{L} \subseteq \binom{X}{2}$ must be a covering of X. Also note that it implies that if $\mathscr{L}$ is a topological lasso for an X-tree T then $\mathscr{L}$ must contain at least $\sum_{v \in V^o(T)} \binom{|V_v|}{2}$ cords and that $\mathscr{L} = \binom{X}{2}$ must hold in case T is the star-tree on X. Finally, note that as the example presented in Fig. 3 shows, the requirement that the two proper edge-weightings in the definition of a topological lasso must be equidistant cannot be dropped. More precisely for $X = \{a,b,c,d\}$ and $\mathscr{L} = \{ab,cd,ad\}$ the X-tree T pictured on the left of that figure (with the indicated edge-weighting ω ignored for the moment) satisfies Properties (C1) and (C2) is not equivalent with the X-tree T' depicted on the right of that figure (again with the indicated edge-weighting ω' ignored where $0 < \varepsilon < 1$). Also ω and ω' are clearly proper and, in the case of ω, also equidistant and (T,ω) and (T',ω') are $\mathscr{L}$-isometric.

Combining Proposition 6.1 and Theorem 7.1, we obtain the following corollary.

Corollary 7.2. *Suppose that T is an X-tree. Then every topological lasso for T must also be an edge-weight lasso for T. Moreover, if T is binary and $\mathscr{L} \subseteq \binom{X}{2}$ then $\mathscr{L}$ is a topological lasso for T if and only if $\mathscr{L}$ is an equidistant lasso for T. In particular, every topological/equidistant lasso for T is also a strong lasso for T in this case.*




FIGURE 3. An example showing that for Theorem 7.1 to hold the "equidistant" requirement in the definition of a topological lasso cannot be dropped (see text for details).

Note however that for $\mathscr{L} \subseteq \binom{X}{2}$ a set of cords and T an X-tree it is possible that $\mathscr{L}$ is simultaneously an edge-weight and a topological lasso for T but T is not binary. We present such an example in the next section.

8. EXAMPLES OF LASSOS

In this section we apply our findings to two types of constructions of sets of cords of X. Both of them were originally introduced in Dress et al (2012b) for the case of edge-weighted, unrooted, phylogenetic trees with leaf set X where an edge-weighting for such a tree is defined as in the case of an X-tree.

Assume for the remainder of this section that T is an X-tree. Then the first example relies on the notion of a circular ordering (of the leaf set) of T (see e.g. Semple and Steel (2003) for further details on such orderings). Following Dress et al (2012b) a *circular ordering (of X) of T* is a cyclic permutation σ of X such that the following holds. There exists a planar embedding of T such that, for every $x \in X$, the leaf that is encountered after x when traversing T in that embedding in say, a counter-clockwise fashion, is the leaf $\sigma(x)$. For example, (a,b,c,d) is a circular ordering of T the $\{a,b,c,d\}$-tree depicted in Figure 3(a).

Let $(x_1,x_2,\ldots,x_n)$ denote a circular ordering of the leaves of T where $n := |X|$ and put $x_{n+1} = x_1$. Then since for every interior vertex v of T there exists some $i \in \{1,\ldots,n\}$ such that v lies on the path from x_i to x_{i+1}, Theorem 6.1 implies that the set

$$\mathscr{L} = \{x_i x_{i+1} : 1 \leq i \leq n\}$$

is an equidistant lasso for T. However, $\mathscr{L}$ is clearly not an equidistant lasso of minimal size. In view of Theorem 7.1, $\mathscr{L}$ is a topological lasso for T if and only if every interior vertex $v \in V^o(T)$ has degree three (except possibly the root ρ_T which might also have degree two) as in that case $G(\mathscr{L},w)$ is a complete graph for all $w \in V^o(T)$. Thus, $\mathscr{L}$ is also a strong lasso for such X-trees. In view of Theorem 5.1, $\mathscr{L}$ is a weak lasso for T if and only if every vertex $v \in V^o(T)^-$ has degree three (except possibly the root ρ_T which might also have degree two) as in that case $G(\mathscr{L},w)$ is rich for all $w \in V^o(T)^-$ and connected for all $w \in V^o(T) - V^o(T)^-$.

The next construction relies on the notion of a bipartition $\{A,B\}$ of X and was introduced in Dress et al (2012b) where it was shown that the set

$$A \vee B := \{ab \in \binom{X}{2} : a \in A \text{ and } b \in B\}$$

is a topological lasso[1] for an unrooted phylogenetic tree T' with leaf set X if and only if, for every 2-subset $\mathfrak{c}$ of X whose elements have that same parent in T', we have that $A\cap\mathfrak{c}\neq\emptyset\neq B\cap\mathfrak{c}$. Note that this implies in particular that for an unrooted phylogenetic tree on X to be topologically lassoed by $A\vee B$, every interior vertex of T can be adjacent with at most two leaves. Thus every pseudo-cherry of T' (defined as in the case of an X-tree) must be a cherry of T.

Defining for an unrooted phylogenetic tree T' on X a set $\mathscr{L}\subseteq\binom{X}{2}$ of cords to be an edge-weight lasso as in the case of an X-tree but again with the requirement "equidistant" on the proper edge-weightings removed, it is not difficult to see that $A\vee B$ is not an edge-weight lasso for T'. Also it is straight forward to see that any two proper edge-weightings ω and ω' for T' such that $D_{(T',\omega)}(a,b)=D_{(T',\omega')}(a,b)$ holds for all $ab\in\mathscr{L}$ must coincide on the interior edges of T' where for a proper edge-weighting α of T' we denote the induced distance on $V(T')$ also by $D_{(T',\alpha)}$.

In the case of X-trees the situation changes in so far that if T is not the star-tree on X and $\{A,B\}$ is such that every pseudo-cherry of T contains elements from both A and B then, in view of Theorem 5.1, $A\vee B$ must be a weak lasso for T and thus, by Corollary 6.2, also an equidistant lasso for T. In view of Theorem 7.1, $A\vee B$ is not a topological lasso for T unless every interior vertex $v\in V^o(T)$ of T is incident with at most two leaves of T and, if v is incident with two leaves, then one is contained in A and the other in B as otherwise $G(A\vee B,v)$ would not be a clique. Since for such X-trees T we have, for all $v\in V^o(T)$, that $G(\mathscr{L},v)$ contains at least one edge it follows that $A\vee B$ is a strong lasso for T.

If T is the star-tree on X then $A\vee B$ is a weak lasso for T. Also, $A\vee B$ is an equidistant lasso for T as $G(\mathscr{L},\rho_T)$ contains at least one edge but it is not a topological lasso for T as $A\vee B\neq\binom{X}{2}$.

9. Conclusion

In the form of investigating when a set of cords of a finite set X of size at least three is an equidistant/topological/weak/strong lasso for an X-tree, we have addressed the topical problem of when a set of partial distances for a set of individuals within a variety uniquely determines a dendogram on those individuals. Such structures are commonly constructed as part of a phenetic clustering step within a genomewide association study to better understand the link between phenotypic and a genotypic variation within a variety. For T an X-tree and $\mathscr{L}\subseteq\binom{X}{2}$ a set of cords, we present characterizations for when $\mathscr{L}$ is an equidistant/weak/tropological lasso for T in terms of the structure of the child-edge graphs associated to the interior vertices of T. As immediate consequences, our characterizations allow us to not only shed light into the problem of when two of the above types of lassos coincide but also into the size of minimum size equidistant/topological/weak lassos.

Despite these encouraging results a number of open questions remain. For example, the characterizations above require knowledge of the structure of T in the form of the child-edge graphs associated to the interior vertices of T. Thus, is it possible to characterize or at least understand lassos without this structural insight into T. A potential candidate for this might be the graph $\Gamma(\mathscr{L})$ associated to $\mathscr{L}$ whose vertex set is X and whose edge set is $\mathscr{L}$. The underlying rational for this is that for $|X|\geq 4$, it was shown in (Dress et al, 2012b, Theorem 1) that for $\mathscr{L}$ to be a topological lasso for an unrooted phylogenetic tree T on X the graph $\Gamma(\mathscr{L})$ has to be connected and for $\mathscr{L}$ to be an edge-weight lasso for T it has

[1]The definition of a topological lasso for an unrooted phylogenetic tree on X is the same as that of a topological lasso for an X-tree but with the requirement dropped that the two proper edge-weightings mentioned in that definition are equidistant.

to be strongly non-bipartite (where a graph G is said to be *strongly non-bipartite* if every connected component of G is not bipartite). Also for $\mathscr{L}$ as constructed in the first example in Section 8 the graph $\Gamma(\mathscr{L})$ is connected.

To overcome the potential loss of information in distance based phylogenetic tree reconstruction, Pachter and Speyer (2004) proposed using *k-dissimilarities*, $k \geq 3$, on X rather than 2-similarities as is the case when reconstructing edge-weighted phylogenetic trees from distances (see also (Felsenstein, 2003, p.176) and Herrmann et al (in press); Warrens (2010); Deza and Rosenberg (2000) and the references therein for recent work on such objects which are sometimes also called k-way similarities, k-way distances, and k-semimetrics). It would be interesting to know what can be said about lassoing and coralling of X-trees within this more general framework.

Acknowledgement A. -A. Popescu thanks the Norwich Research Park (NRP) for support.

REFERENCES

De Soete G (1984) Ultrametric tree representations of incomplete dissimilarity data. Journal of Classification 1:235–242

Deza MM, Rosenberg IG (2000) n-semimetrics. European J Combin 21(6):797–806

Diestel R (Electronic Edition 2005) Graph Theory. Springer Verlag, Heidelberg, New York

Dress A, Huber K, Koolen J, Moulton V, Spillner A (2012a) Basic Phylogenetic Combinatorics. Cambridge University Press, Cambridge, UK

Dress AWM, Huber KT, Steel M (2012b) 'Lassoing' a phylogenetic tree I: basic properties, shellings and covers. Journal of Mathematical Biology 65:77–105

Felsenstein F (2003) Inferring Phylogenies. Sinauer Associates

Harper A, Trick M, Higgins J, Fraser F, Clissold L, Wells R, Hattori C, Werner P, Bancroft I (in press) Associative transcriptomics of traits in the polyploid crop species brassica napus. Nature Biotechnology

Herrmann S, Huber KT, Moulton V, Spillner A (in press) Recognizing treelike k-dissimilarities. Journal of Classification

Muir WM, Wong GKS, Zhang Y, Wang J, Groenen MAM, Crooijmans RPMA, Megens HJ, Zhang H, Okimoto R, Vereijken A, Jungerius A, Albers GAA, Lawley CT, Delany ME, MacEachern S, Cheng HH (2008) Genome-wide assessment of worldwide chicken SNP genetic diversity indicates significant absence of rare alleles in commercial breeds. Proceedings of the National Academy of Sciences 105(45):17,312–17,317

Pachter L, Speyer D (2004) Reconstructing trees from subtree weights. Applied Mathematics Letters 17:615–621


Semple C, Steel M (2003) Phylogenetics, Oxford Lecture Series in Mathematics and its Applications, vol 24. Oxford University Press, Oxford, UK


Warrens MJ (2010) n-way metrics. J Classification 27(2):173–190


Zhang Z, Ersoz E, Lai CQ, Todhunter RJ, Tiwari HK, Gore MA, Bradbury PJ, Yu J, Arnett DK, Ordovas JM, Buckler E (2010) Mixed linear model approach adapted for genome-wide association studies. Nature Genetics 42(4):355–60


SCHOOL OF COMPUTING SCIENCES, UNIVERSITY OF EAST ANGLIA, NORWICH, NR4 7TJ, UK
E-mail address: katharina.huber@cmp.uea.ac.uk


SCHOOL OF COMPUTING SCIENCES, UNIVERSITY OF EAST ANGLIA, NORWICH, NR4 7TJ, UK
E-mail address: Andrei-Alin.Popescu@uea.ac.uk


T :
ρ_T
1
1
2
2
2
2
2
a
b
c
d
e
T' :
$\rho_{T'}$
1
1
1
2
1
1
2
2
a
c
b
d
e



T
1
1
1
1
1
1
1
1
a
b
c
d
e


T'
0.5
0.5
1
3
2.5
1
1
2
b
c
d
a
e


ρ_T
$T:$
1
2
2
1
1
1
1
a b c d e f
$\rho_{T'}$
$T':$
1
2
1
1
1
1
1
a b c d f e



$a)$



$b)$